

Planting Rate and Date

Rates are based on seed of normal size and good quality and normal seedbed. Actual rates used will vary widely, depending on seed cost, desired stand, expected mortality, emerging ability, seed weight, seed germination, seedbed condition, depth of planting and planting equipment.

Crop	Bushel Weight (Pounds) ¹	Seeds / Pound (Number)	Rate / Acre (Pounds)	Rate (Seeds)	Planting Date
Barley	48	14,300	85	28 / sq. ft.	Early spring
Corn	56	—		33,000 / acre	April 15 / May 5
Fieldbean					
Black turtle soup	60	2,300	45	105,000 / acre	May 20 / June 15
Great northern	60	1,000	100	90,000 / acre	May 20 / June 15
Kidney	60	900	90-115	90,000 / acre	May 20 / June 15
Navy	60	2,500	42	105,000 / acre	May 20 / June 15
Navy, rows 6 to 14 in.	60		60	150,000 / acre	May 20 / June 15
Pinto	60	1,300	80	90,000 / acre	May 20 / June 15
Small red	60	1,400	75	100,000 / acre	May 20 / June 15
Small white	60	3,000	35	105,000 / acre	May 20 / June 15
Flax	56	88,000	42	85 / sq. ft.	April 15 / May 15
Forage grasses, perennial					
Bromegrass alone	14	136,000	16	50 / sq. ft.	Early spring or late summer
Bromegrass in mixtures			5	15 / sq. ft.	Use date for legumes
Orchardgrass, alone	14	653,000	10	150 / sq. ft.	Early spring or late summer
Orchardgrass, in mixtures			3	45 / sq. ft.	Use date for legumes
Reed canarygrass alone	46	526,000	7	85 / sq. ft.	Early spring or late summer
Reed canarygrass, in mixtures			5	60 / sq. ft.	Use date for legumes
Tall fescue, alone	25	229,000	15	75 / sq. ft.	Early spring or summer
Tall fescue, in mixtures			5	20 / sq. ft.	Use date for legumes
Timothy	45	1,234,000	3	85 / sq. ft.	Use date for legumes
Forage legumes, perennial					
Alfalfa alone	60	220,000	13	65 / sq. ft.	Late April-early May / Late June-early August
Alfalfa with grass			5 to 10	25 to 50 / sq. ft.	Late April-early May / Late June-early August
Alsike clover	60	653,000	2	30 / sq. ft.	Early spring to August 10
Birdsfoot trefoil alone	60	372,000	8	70 / sq. ft.	Early spring or summer
Birdsfoot trefoil in mixtures			6	50 / sq. ft.	Early spring or summer
Cicer milkvetch	60	122,000	18	50 / sq. ft.	Early spring or summer
Ladino clover	60	784,000	1	18 / sq. ft.	Early spring to August 10
Red clover alone	60	272,000	9	55 / sq. ft.	Early spring to September 1
Red clover with grass			5	30 / sq. ft.	Use date for legumes
Oat	32	16,200	80	28 / sq. ft.	Early spring
Rye	56	18,200	60	25 / sq. ft.	September 1
Sorghum, rows 18 to 40 in.	56	15,000	10	150,000 / acre	May 20 to June 5 for grain
Sorghum, rows 6 to 14 in.			15	5 / sq. ft.	
Soybean, 7-in. rows	60	2,800	56	2 / ft. of row	May 1 to May 10
10-in. rows				3 / ft. of row	
20-in. rows				6 / ft. of row	
22-in. rows				7 / ft. of row	
30-in. rows				9 / ft. of row	
Sunflower, nonoilseed	24	4,300	4	17,000 / acre	May 1—June 15
Sunflower, oilseed	27	7,700	3	23,000 / acre	May 1—June 15
Wheat, durum	60	12,100	90	25 / sq. ft.	Early spring
Wheat, hard red spring ²	60	14,000	113	28 / sq. ft.	Early spring
Wheat, hard red winter	60	14,500	75+	25 / sq. ft.	August 20 / September 20
Other Crops					
Annual canarygrass	50	58,000	30	40 / sq. ft.	Early spring
Buckwheat	48	14,900	50	17 / sq. ft.	June 15 / July 20
Canola, <i>B. napus</i>	50	80,000 to 160,000	3 to 5	6 to 9	Early spring
Crambe	22	65,000	15	23 / sq. ft.	Late April / early May
Fieldpea	60	2,300	180	9 / sq. ft.	Early spring
Fieldpea with 1½ to 2 bu. oat			70	4 / sq. ft.	Early spring
Fababean, medium size	60	1,300	180	5 / sq. ft.	Early spring
Fababean, with 2 bu. oat			60	2 / sq. ft.	Early spring
Lentil, small	60	15,600	55	20 / sq. ft.	Early spring
Millet, foxtail	48	218,000	15	75 / sq. ft.	June 15 / July 15
Millet, proso	56	65,000	20	30 / sq. ft.	June 15 / July 15
Sudangrass, rows 6 to 14 in.	40	44,000	25	25 / sq. ft.	May 20 / June 10
Sweetclover	60	240,000	10	55 / sq. ft.	Early spring
Wildrice (wet)	25	7,900	35	6 / sq. ft.	Late fall

¹ U.S. legal bushel weight or, if not established, the weight most widely accepted. ² See wheat section for best way to calculate hard red spring wheat planting rate.