

# BROMEGRASS

Bromegrass is generally grown for hay in mixture with alfalfa, or is used as pasture in mixture with other grasses and legumes. Varieties can be classed as southern, intermediate and northern types. Varieties of the southern type may

not be higher yielding but are generally less susceptible to leaf diseases and earlier in maturity than northern types. All varieties are winterhardy. Some stand losses may occur when bromegrass is managed under three- and four-cut alfalfa harvest systems. Overgrazing also will reduce stands.

Varieties are generally evaluated in pure stands at Minnesota experiment stations with a three-cut harvest system. Nitrogen was applied at all locations in early spring and after the first two harvests at a rate of 50 pounds per acre. Average dry matter yields were very high in 1998-99 and few differences were found among performance of the varieties. Stand losses on all entries included in the table were noted in Rosemount after the first harvest in 1999, except for the variety York.

***Dry-matter yield, in tons dry matter per acre, of smooth bromegrass seeded at two locations.***

Variety	Rosemount	Morris
	1998-2000	1999-2001
Alpha	4.8	4.6
Badger	4.4	4.5
Bounty	4.8	4.7
Lincoln	4.5	–
York	5.2	4.7
LSD 5%	0.3	NS

***Bromegrass  
Planting Rate and Date***

Bushel Weight, Pounds .....	14
Seeds/Pound .....	136,000
Planting Rate, Pounds/Acre	
Alone .....	16
In Mixtures .....	5
Planting Rate, Seeds Sq.Ft.	
Alone .....	50
In Mixtures .....	15
Planting Date .....	Early Spring or Late Summer