

# Minnesota

## VARIETAL TRIALS RESULTS

---

To help growers select varieties best adapted to a specific area, the Minnesota Agricultural Experiment Station (MAES) compares varieties in trial plots at St. Paul, Becker, Crookston, Grand Rapids, Lamberton, Morris and Waseca, and in farmers' fields. Crop varieties are grown in replicated plots at each location, and factors affecting their yield and characteristics are as nearly the same as possible for all varieties at each location.

Not all crop varieties included in previous editions of Minnesota Varietal Trials are included in this 2008 edition. If you have a question about such a crop, contact the author(s) of the related crop section at 612-625-2740 or go to the MAES web site at [www.maes.umn.edu](http://www.maes.umn.edu). Click on Commodity/Crop Lines, then on Varietal Trials, which will display a crop listing.

### **Certified Seed**

Use of certified seed is suggested, but certification in itself does not imply recommendation. Registered and certified seed of most varieties described in this report can be purchased from seed dealers or grower-members of the Minnesota Crop Improvement Association (MCIA). You can find sources of certified and registered seed at the MCIA web site, [www.mcia.org](http://www.mcia.org).

### **Interpreting the Tables**

The LSD (least significant difference) numbers beneath yield columns in tables are statistical measures of variability within trials. The LSD is used to determine whether the difference between two yields is due to a genetic difference in the varieties or to other causes, such as environmental variability.

If the yield difference between two varieties equals or exceeds the LSD value for the yield column, the higher-yielding variety probably was superior in yield. If the difference is less than the LSD the yield difference probably was due to environmental factors. An "NS" notation in a column indicates no significant difference for that characteristic. The relative maturities of varieties are variously indicated in the tables as date of maturity; date of heading or blooming; days to maturity, heading or blooming; or moisture percentage at harvest.

These varietal trials are not designed for crop (species) comparisons; crops are grown on dif-

ferent fields and with different management. The data should be used only to compare varieties within a table.

### **Abbreviations**

To save space in variety descriptions and some other listings, "agricultural experiment station" often is abbreviated as AES.

### **Cooperating Researchers**

Ruth Dill-Macky, Senyu Chen, Jim Kurle and Dean Malvick, Department of Plant Pathology; Y Jin and Jim Kolmer, USDA-ARS Cereal Disease Laboratory; and Gary Hareland, USDA-ARS Wheat Quality Laboratory, Fargo.

### **Publication Staff and Information**

**Coordination:** Jennifer Obst

**Supervisor:** Lee Hardman

**Photography:** Dave Hansen

Information in this publication is published under authority granted by the Hatch Act of 1887 to conduct performance trials of farm crops and interpret data to the public.

Permission is granted to reproduce tables only in their entirety, without rearrangement, manipulation or reinterpretation. Permission also is granted to reproduce a maturity group sub-table, provided that complete table headings and table notes are included. Reproductions of any material from this publication should credit the Minnesota Agricultural Experiment Station as its source.

In accordance with the Americans with Disabilities Act, information in *Minnesota Varietal Trials Results* is available in alternative forms upon request. Contact the University of Minnesota Distribution Center at 612-625-8173.

The University of Minnesota Agricultural Experiment Station is an equal opportunity employer.

Copyright © 2007, Regents of the University of Minnesota, Minneapolis, MN 55455.

Minnesota Agricultural  
Experiment Station

---

UNIVERSITY OF MINNESOTA