

# TIMOTHY

Timothy is adapted throughout Minnesota for use in hay and pasture mixes. When timothy is the major component in hay, its stage of maturity affects both yield and quality. Harvesting timothy at early heading is the preferred time. Timothy produces the majority of its forage at the first harvest.

Varieties of timothy differ in maturity so care should be taken in choosing ones that fit the management requirements of the crop and mixture. Early varieties are best adapted to a three-cut system with alfalfa.

Intermediate to late-maturing varieties should not be harvested more than twice during the growing season. Appropriately selected timothy varieties are compatible with red clover and birdsfoot trefoil in mixtures for hay production.

Varieties in the experiment station timothy trials were established in pure stands in 1992 at Rosemount and Morris and again at Grand Rapids in 1993. Nitrogen was applied at all locations in the early spring and after each harvest at a rate of 40 to 50 pounds per acre.

Early-maturing varieties of timothy had greater forage production than the late maturing varieties at all locations over all harvest years. At Morris and Rosemount the yields of timothy were exceptionally high in 1993 and 1994. These results may be partially attributed to mild winters, and abundant rainfall and cool temperatures during the growing seasons. Timothy is normally less persistent than other cool-season grasses, such as reed canarygrass.

## *Timothy Planting Rate and Date*

Bushel Weight, Pounds .....	45
Seeds/Pound.....	1,234,000
Planting Rate, Pounds/Acre	
In Mixtures.....	3
Planting Rate, Seeds Sq.Ft.	
In Mixtures.....	85
Planting Date	
In Mixtures.....	Use Date for Legume

## *Dry matter yield, tons dry matter per acre, of timothy seeded at three locations.*

Variety	Grand Rapids 1994-96	Rosemount 1993-1995	Morris 1993-1996	Mean
<b>Early-Intermediate maturity:</b>				
Climax	3.6	3.8	4.0	3.8
Comtal	3.4	3.7	–	3.6
Goliath	3.4	3.4	–	3.4
Timfor	3.5	3.8	–	3.7
Toro	3.7	3.9	–	3.8
<b>Late maturity:</b>				
Heidemij	3.5	3.0	3.5	3.3
Hokusen	3.3	3.4	3.6	3.4
LSD 5%	0.4	0.4	NS	